

$SU(3) \times SU(2) \times U(1)$ Chiral Models from Intersecting D4-/D5-Branes

Hironobu KATAOKA^{1*} and Masafumi SHIMOJO^{2**}

¹ *Department of Physics, Hyogo University of Education, Yashiro-cho, Hyogo 673-1494, Japan,*

² *Department of Electronics and Information Technology, Fukui National College of Technology, Sabae 916-8507, Japan*

(Received December 28, 2001)

We clarify RR tadpole cancellation conditions for intersecting D4-/D5-branes. We determine all of the D4-brane models that have D=4 three-generation chiral fermions with $SU(3) \times SU(2) \times U(1)^n$ symmetries. For the D5-brane case, we present a solution to the conditions that gives exactly the matter content of the standard model with $U(1)$ anomalies.

Intersecting Dp-branes are useful tools to obtain chiral fermions existing on their intersections.¹⁾ Although configurations of type IIB D6-branes wrapped on $\mathbf{T}^6$ give rise to identically the matter content of the standard model,²⁾ the D6-brane models do not solve the fine-tuning problem caused by its breakdown of supersymmetry. Contrastingly, this difficulty is avoided by using D4-/D5-branes wrapped on $\mathbf{T}^{2d} \times \mathbf{R}^{6-2d}/Z_N$ through reduction of the string scale with a very large volume of the Z_N orbifold. Intersecting D4-/D5-branes in both oriented and unoriented theories, we construct D=4 chiral models whose matter content transforms as $(3, 2)_{\frac{1}{6}} + (\bar{3}, 1)_{-\frac{2}{3}} + (\bar{3}, 1)_{\frac{1}{3}}$ under $SU(3) \times SU(2) \times U(1)_Y$.

We start with the definition of the spacetime geometry in the oriented theory. The compactified six dimensional space is the product of d rectangular two-tori $\mathbf{T}_I^2$ ($1 \leq I \leq d \leq 2$) and a $\mathbf{R}^{6-2d}/Z_N$ orbifold. The space is parameterized by the complex coordinates $Y_I = X_{2I+2} + iX_{2I+3}$ with radii R_{2I+2} and R_{2I+3} for $I = 1, 2, 3$.

Let us consider K different stacks, each of which is a set of N_a coincident D4-/D5-branes. They are labeled by the index a ($a = 1, \dots, K$) and wrapped around a 1-cycle denoted by $(n_a^{(I)}, m_a^{(I)})$ on each of the d two-tori, where $I = 1$ for D4-branes and $I = 1, 2$ for D5-branes. When the a -th stack and the b -th stack intersect, we refer to their intersections as an a - b sector. The intersection number of the a - b sector is given by

$$I_{ab} = \prod_{I=1}^d I_{ab}^{(I)} = \prod_{I=1}^d (n_a^{(I)} m_b^{(I)} - n_b^{(I)} m_a^{(I)}). \quad (1)$$

The Z_N action is realized by powers of the twist generator represented by a shift vector v in the space of the $SO(6-2d)$ Cartan subalgebra. In the configurations with intersecting D4-branes, modular invariance conditions for one-loop amplitudes

* E-mail: kataoka@yukawa.kyoto-u.ac.jp

** E-mail: shimo0@ei.fukui-nct.ac.jp

restrict the form of the twist vector $v = (v_1, v_2, v_3)$ to $(0, 1, b_2)/N$, where b_2 is an odd integer. For the case of D5-branes, $v = (0, 0, 2/N)$.

Since N_a coincident D-branes have $U(N_a)$ gauge factors in their world-volume, open strings stretched between the a -th stack and the b -th stack give rise to states in the a - b sector that belong to bifundamental representations of $U(N_a) \times U(N_b)$. The gauge degrees of freedom are expressed by the Chan-Paton (CP) factors as $|ij\rangle$, where i and j correspond to a D-brane in the a -th and the b -th stack, respectively. We express the embedding of the Z_N action θ^k using the unitary matrix γ_k . We diagonalize γ_k so that

$$\gamma_k = \text{diag}(\zeta^{l_1} \mathbf{1}_{N_1}, \zeta^{l_2} \mathbf{1}_{N_2}, \dots, \zeta^{l_K} \mathbf{1}_{N_K}), \quad (2)$$

where $\zeta = e^{2\pi i \frac{1}{N}}$ and l_a are integers. Then the N_a D-branes in the a -th stack have the same CP phase, ζ^{l_a} .

The RR tadpole cancellation conditions^{1) 3)} are

$$\sum_{a=1}^K N_a \prod_{I=1}^d n_a^{(I)} = \sum_{a=1}^K N_a \prod_{I=1}^d m_a^{(I)} = 0, \quad (3)$$

$$\sum_{k=1}^{N-1} \left(\prod_{J=d+1}^3 |\sin \pi k v_J| \right) \left(\sum_{a=1}^K \left(\prod_{I=1}^d n_a^{(I)} \right) \text{Tr}_a \gamma_k \right)^2 = 0, \quad (4)$$

$$\sum_{k=1}^{N-1} \left(\prod_{J=d+1}^3 |\sin \pi k v_J| \right) \left(\sum_{a=1}^K \left(\prod_{I=1}^d m_a^{(I)} \right) \text{Tr}_a \gamma_k \right)^2 = 0, \quad (5)$$

where Tr_a represents the trace over the CP factor on D-branes that belong to the a -th stack. For each shift vector v of Z_N , we can obtain solutions for 1-cycles $(n_a^{(I)}, m_a^{(I)})$ of the a -th stack and CP phases ζ^{l_a} . The solution produces an open string spectrum that is invariant under the Z_N action.⁴⁾

For the D4-brane case, left-handed fermions in the a - b sector are given in bifundamental representations under $U(N_a) \times U(N_b)$ as¹⁾

$$I_{ab} \{ (N_a^i, \overline{N_b}^{i - \frac{1-b_2}{2}}) + (N_a^i, \overline{N_b}^{i + \frac{1-b_2}{2}}) + (\overline{N_a}^i, N_b^{i - \frac{1+b_2}{2}}) + (\overline{N_a}^i, N_b^{i + \frac{1+b_2}{2}}) \}, \quad (6)$$

where the symbol N_a^i expresses the meaning that this is the representation N_a of $U(N_a)$ and the CP phase of the a -th stack is ζ^i . A negative value of I_{ab} indicates a positive multiplicity of the field that transforms as the complex conjugate of each bifundamental representation. To obtain the three (3,2) representations, we require $N_1 = 3$ and $N_2 = 2$ stacks and the intersection number $I_{12} = 3$. The two types of three-generation right-handed quarks require $N_3 = 1$ and $N_4 = 1$ stacks and intersection numbers $I_{13} = \pm 3$ and $I_{14} = \pm 3$. One more stack $N_5 = 1$ is necessary for the consistency conditions (3)–(5) to be satisfied. We have investigated all of these configurations and found two solutions, given in Table I.

Since the gauge symmetry $U(1)$ of the stack $N_a = 1$ with 1-cycle $(n_a, 0)$ extends to $U(1)^{|n_a|}$,¹⁾ the gauge symmetry of the models is $U(3) \times U(2) \times U(1)^5 = SU(3) \times$

Table I. D4-brane 1-cycles and CP phases giving rise to three generations of quarks.

number	Z_N orbifold	b_2	cycle				
			CP phase				
			$N_1 = 3$	$N_2 = 2$	$N_3 = 1$	$N_4 = 1$	$N_5 = 1$
D4-1	$N \geq 3$	$b_2 \neq -1$	$(1, 0)$ 1	$(n_2, 3)$ $\zeta^{\frac{b_2-1}{2}}$	$(n_3, -3)$ $\zeta^{\frac{b_2-1}{2}}$	$(-2n_2 - n_3, -3)$ $\zeta^{\frac{b_2-1}{2}}$	$(-3, 0)$ 1
D4-2	$N \geq 3$	$b_2 = -1$	$(1, 0)$ 1	$(n_2, 3)$ $\zeta^{\frac{b_2-1}{2}}$	$(n_3, -3)$ $\zeta^{\frac{b_2-1}{2}}$	$(-2n_2 - n_3, -3)$ $\zeta^{\frac{b_2-1}{2}}$	$(-3, 0)$ 1

$SU(2) \times U(1)^7$. The fermion spectrum under the symmetry given by D4-1 in Table I is as follows:

$$3(3, 2)_{(1, -1, 0, 0, 0^3)} + 3(\bar{3}, 1)_{(-1, 0, 1, 0, 0^3)} + 3(\bar{3}, 1)_{(-1, 0, 0, 1, 0^3)} \\ + 3(1, 2)_{(0, 1, 0, 0, \underline{-1, 0, 0})} + 3(1, 1)_{(0, 0, -1, 0, \underline{1, 0, 0})} + 3(1, 1)_{(0, 0, 0, -1, \underline{1, 0, 0})}, \quad (7)$$

where the underlines indicate permutation of indices. Computing the mixed anomalies which need to be cancelled by the Green-Schwartz mechanism, we find six non-anomalous $U(1)$ linear combinations. They include the suitable hypercharge

$$Q_Y = -\frac{1}{6}Q_2 + \frac{1}{3}Q_3 - \frac{2}{3}Q_4 + \frac{1}{3}(Q_5^{(1)} + Q_5^{(2)}) - \frac{2}{3}Q_5^{(3)}, \quad (8)$$

where Q_a is the generator of the a -th $U(1)$. With this hypercharge, the fermion spectrum is

$$3(3, 2)_{\frac{1}{6}} + 3(\bar{3}, 1)_{\frac{1}{3}} + 3(\bar{3}, 1)_{-\frac{2}{3}} + 6(1, 2)_{-\frac{1}{2}} + 3(1, 2)_{\frac{1}{2}} \\ + 6(1, 1)_1 + 9(1, 1)_0 + 3(1, 1)_{-1}. \quad (9)$$

When we set $n_2 + n_3 = 1$, D4-2 in Table I gives the following fermion spectrum:

$$3(3, 2)_{(1, -1, 0, 0, 0^3)} + 3(\bar{3}, 1)_{(-1, 0, 1, 0, 0^3)} + 3(\bar{3}, 1)_{(-1, 0, 0, 1, 0^3)} \\ + 3(1, 2)_{(0, 1, 0, 0, \underline{-1, 0, 0})} + 3(1, 1)_{(0, 0, -1, 0, \underline{1, 0, 0})} + 3(1, 1)_{(0, 0, 0, -1, \underline{1, 0, 0})} \\ + 6(1, 2)_{(0, 1, -1, 0, 0^3)} + 6(1, 2)_{(0, -1, 0, 1, 0^3)} + 12(1, 1)_{(0, 0, 1, -1, 0^3)}. \quad (10)$$

We choose the hypercharge from anomaly-free realizations of $U(1)$ as

$$Q_Y = \frac{1}{3}Q_1 + \frac{1}{6}Q_2 + \frac{2}{3}Q_3 - \frac{1}{3}Q_4 - \frac{1}{3}(Q_5^{(1)} + Q_5^{(2)} + Q_5^{(3)}). \quad (11)$$

With this hypercharge, the spectrum (10) become

$$3(3, 2)_{\frac{1}{6}} + 3(\bar{3}, 1)_{\frac{1}{3}} + 3(\bar{3}, 1)_{-\frac{2}{3}} + 12(1, 2)_{-\frac{1}{2}} + 9(1, 2)_{\frac{1}{2}} \\ + 12(1, 1)_1 + 9(1, 1)_0 + 9(1, 1)_{-1}. \quad (12)$$

When we set $n_2 + n_3 = -1$, we get the same spectrum (12) by exchanging Q_3 and Q_4 in the hypercharge (11). For the case $n_2 + n_3 \neq \pm 1$, we have not been able to obtain the solution with the correct hypercharge Q_Y .

When the a -th stack and b -th stack have the same CP phase, there are I_{ab} tachyon fields in the a - b sector.¹⁾ In these models, the 2-3 and 2-4 sectors give rise to tachyon fields that transform as the $SU(2)$ doublet. This indicates that the configurations of branes that have $U(2)$ and $U(1)$ symmetries are unstable. This instability suggests a stringy Higgs mechanism of electroweak symmetry breaking.¹⁾

For the D5-brane case, the fact that there are many wrapping numbers ($n_a^{(I)}$, $m_a^{(I)}$) with ($I = 1, 2$) makes it difficult to obtain general solutions. For this reason, in this paper, we content ourselves with a single example. We will return to this problem at the end of the investigation of the unoriented theory.

We now discuss the unoriented theory. We introduce an orientifold group written as $Z_N + \Omega R Z_N$, where Ω is the world sheet parity. R transforms Y_I to $\overline{Y_I}$ for $1 \leq I \leq d$ and Y_I to $-Y_I$ for $d+1 \leq I \leq 3$. The space-time is expressed as

$$\frac{\mathcal{M}_4 \times \prod_{I=1}^d \mathbf{T}_I^2 \times (\prod_{I=1+d}^3 \mathbf{T}_I^2 / Z_N)}{\Omega R}.$$

We express the ΩR action on the CP factors using unitary matrices $\gamma_{\Omega R}$. Since $\Omega R \theta^k (\Omega R)^{-1} = \theta^{2k}$, we have

$$\gamma_{\Omega R k} = \gamma_k \gamma_{\Omega R} = \pm \gamma_{2k} \gamma_{\Omega R k}^T. \quad (13)$$

This leads to $\gamma_{\Omega R} = \pm \gamma_{\Omega R}^T$. In the following equations, the upper and lower sign correspond to the symmetric and antisymmetric $\gamma_{\Omega R}$, respectively.

In addition to the a - b sector, there are intersections of the a -th stack and mirror branes of the b -th stack, which we refer to as a - b^* sector. The intersection number I_{ab^*} is obtained by changing $m_b^{(I)}$ to $-m_b^{(I)}$ in the expression (1).

The Z_N action with even N contains an order 2 element, so that the orientifold group contains an element that does not vary Y_I ($I \geq d+1$). Then there will be D8-branes in the type IIA D4-brane theory. Since the concept of intersecting D-branes involves use of the same dimensional D-branes, we restrict ourselves to the case that the order N of Z_N is odd.

Tadpole divergences come from Klein bottle and Möbius strip amplitudes as well as cylinder amplitudes.³⁾ The RR tadpole cancellation conditions common to configurations with intersecting D4-branes and configurations with D5-branes are given by

$$\sum_{a=1}^K N_a \prod_{I=1}^d n_a^{(I)} = \pm 16, \quad (14)$$

$$\sum_{a=1}^K N_a \prod_{I=1}^d m_a^{(I)} = 0, \quad (15)$$

$$\sum_{a=1}^K \left(\prod_{I=1}^d m_a^{(I)} \right) \text{Tr}_a \gamma_k = 0 \quad \text{for } k = 1, \dots, N-1. \quad (16)$$

There is one more condition on the product of the cycles $n_a^{(I)}$ and the trace of γ_k .

When we consider a Z_3 orbifold, it is given by

$$\sum_{a=1}^K n_a^{(1)} \text{Tr}_a \gamma_k = \pm 4 \quad \text{for } k = 1, 2, \quad (17)$$

for the D4-brane case and

$$\sum_{a=1}^K n_a^{(1)} n_a^{(2)} \text{Tr}_a \gamma_k = \mp 8 \quad \text{for } k = 1, 2, \quad (18)$$

for the D5-brane case. For other Z_N models with odd N , the condition is expressed by a linear summation of $\prod_{J=d+1}^3 \sin \pi k v_J$ and $\prod_{J=d+1}^3 \sin 2\pi k v_J$ over $k = 1, \dots, N-1$ similar to (4), and we could not find any solution to 1-cycles and γ matrices.

In order to satisfy the conditions (14) and (17) for unoriented D4-brane configurations giving a standard-like model, we must introduce many $U(1)$ stacks or an $N_2 = 2$ stack that has a 1-cycle $n_2^{(1)} \neq 1$. Some of these $U(1)$ stacks and the $N_1 = 3$ stack always have intersections that lead to non-standard $(3, 1)$ or $(\bar{3}, 1)$ representations under $SU(3) \times SU(2)$. The 1-cycles $n_2^{(1)} \neq 1$ in the unoriented theory produce fields that transform as a three-dimensional representation under $U(2)$.²⁾ Thus we were not able to obtain the matter content of the standard model.

For models with intersecting D5-branes, left-handed fermions in the a - b sector and a - b^* sector are given by^{1) 2)}

$$I_{ab}((N_a^i, \overline{N}_b^{i+1}) + (\overline{N}_a^i, N_b^{i-1})), \quad (19)$$

$$I_{ab^*}((N_a^i, N_b^{-i-1}) + (\overline{N}_a^i, \overline{N}_b^{-i+1})), \quad (20)$$

under $U(N_a) \times U(N_b)$.

The bifundamental representations of (20) for the a - a^* sector change to $N_a^i \wedge N_a^{-i-1}$ and $\overline{N}_a^i \wedge \overline{N}_a^{-i+1}$ under $U(N_a)$. On-orientifold intersections of the a - a^* sector give $4m_a^{(1)} m_a^{(2)}$ fermions in the representation of either an antisymmetric or symmetric tensor, depending on whether $\gamma_{\Omega R}$ is antisymmetric or symmetric. Off-orientifold intersections of the a - a^* sector produce $2m_a^{(1)} m_a^{(2)} (n_a^{(1)} n_a^{(2)} - 1)$ symmetric and antisymmetric representations. We require that any brane stack satisfy $m_a^{(1)} m_a^{(2)} = 0$ to avoid the appearance of exotic quantum numbers and to satisfy the RR tadpole cancellation conditions (15) and (16). Adding D5-branes with $m^{(1)} = m^{(2)} = 0$ will satisfy the conditions (14) and (18) without modifying the fermionic matter content.

Since $I_{12} = -I_{12^*}$ under 1-cycles for which $m_a^{(1)} m_a^{(2)} = 0$, we must set $I_{12} = \pm 3$ and choose CP factors yielding no fermions in the 1-2* sector to get $3(3, 2)$ representations of $U(3) \times U(2)$. This configuration makes it impossible to obtain just the standard model spectrum without $U(1)$ - $U(1)$ - $U(1)$ anomalies.²⁾ An example is given in Table II for the case in which the orbifold group is Z_3 .

The spectrum under the gauge symmetry $SU(3) \times SU(2) \times U(1)$ ⁵ is given by

$$\begin{aligned} & 3(3, 2)_{(1, -1, 0, 0, 0)} + 3(\bar{3}, 1)_{(-1, 0, 1, 0, 0)} + 3(\bar{3}, 1)_{(-1, 0, 0, 1, 0)} \\ & + 3(1, 2)_{(0, 1, 0, 0, -1)} + 3(1, 1)_{(0, 0, -1, 0, 1)} + 3(1, 1)_{(0, 0, 0, -1, 1)}. \end{aligned} \quad (21)$$

Table II. Example of D5-brane 1-cycles and CP phases giving rise to three generations of quarks. The orbifold group is Z_3 and the parameter ζ is $e^{2\pi i/3}$.

N_a	$(n_a^{(1)}, m_a^{(1)})$	$(n_a^{(2)}, m_a^{(2)})$	CP phase
$N_1 = 3$	$(n_1^{(1)}, 0)$	$(n_1^{(2)}, 3/n_1^{(1)})$	ζ^2
$N_2 = 2$	$(n_2^{(1)}, m_2^{(1)})$	$(1/m_2^{(1)}, 0)$	ζ
$N_3 = 1$	$(n_3^{(1)}, m_3^{(1)})$	$(-1/m_3^{(1)}, 0)$	ζ
$N_4 = 1$	$(n_4^{(1)}, m_4^{(1)})$	$(-1/m_4^{(1)}, 0)$	ζ
$N_5 = 1$	$(n_5^{(1)}, 0)$	$(n_5^{(2)}, -3/n_5^{(1)})$	ζ^2

Although the spectrum has a $U(1)$ - $U(1)$ - $U(1)$ anomaly and G^2 - $U(1)$ mixed anomalies that must be removed by some mechanism, two anomaly-free $U(1)$ linear combinations exist. We can define the hypercharge as

$$Y = \frac{1}{6}Q_1 + \frac{1}{2}(Q_3 - Q_4 + Q_5). \quad (22)$$

and obtain identically the standard model spectrum under $SU(3) \times SU(2) \times U(1)_Y$ gauge symmetry with three generations of right-handed neutrino.

We now consider the construction of D5-branes in the oriented theory. In this case, any matter fermions (21) for the unoriented theory are obtained from the a - b sector, not from the a - b^* sector. Adding an appropriate number of $U(1)$ branes to the stacks in Table II, we obtain D5-branes for the oriented theory that satisfy the conditions on the 1-cycles and CP phases (3)–(5).

Configurations with intersecting D-branes cause the breakdown of supersymmetry. Then the string scale must be close to the weak scale to avoid the fine-tuning problem. The four dimensional Plank mass M_p and string scale M_s are related as $M_p \approx 4\pi^2 M_s^4 \sqrt{V_T V_{Z_N}} / \lambda_{II}$, where λ_{II} is the Type II string coupling, V_T is the volume of the tori, and V_{Z_N} is the volume of the orbifold. While a very large value of V_T leads to small Yukawa and gauge couplings, we can give a very large volume to the transverse Z_N orbifold. For models of D5-branes, for example, we can set $M_s \approx 1 \sim 10$ TeV and $V_T \sim 1/M_s^2$. In order to obtain very large value of the Plank mass, we should choose $\sqrt{V_{Z_N}} \approx 10^9 - 10^{11} \text{ (GeV)}^{-1}$, i.e., $10^{-2} - 10^{-4} \text{ cm}$.

In this paper, we have not investigated all of the standard-like spectra that accompany intersecting D5-branes. In particular, unoriented models may be obtained from 1-cycles for which $m_a^{(1)} m_a^{(2)} \neq 0$, $n_a^{(1)} n_a^{(2)} = 1$. In such models, the a - a^* sector may produce the antisymmetric tensors of $N_a \wedge N_a$ under $U(N_a)$ for $N_a = 2, 3$ without symmetric tensors that are not present in the matter content of the standard model.

-
- [1] G. Aldazabal, S. Franco, L. E. Iáñez and R. Rabadan, A. M. Uranga, hep-ph/0011132, hep-th/0011073.
 - [2] L. E. Ibáñez, F. Marchesano, R. Rabadan, hep-th/0105155.
 - [3] R. Blumenhagen, L. Görlich, B. K rs and D. L st, hep-th/0007024, G. Aldazabal, A. Font, L. E. Ib  nez and G. Violer , hep-th/9804026, S. Ishihara, H. Kataoka and H. Sato, Phys. Rev. D60 (1999), 126005.
 - [4] M. R. Douglas and G. Moore, hep-th/9603167.